

AN IMPORTANT ALERT TO KEEP WITH YOU

In case of a medical emergency, healthcare providers may ask for your medical history to help choose the appropriate medication for you. This Patient Alert Card will let healthcare providers know that you are being treated with SUBLOCADE® (buprenorphine extended-release).

COMPLETE THE CARD, CUT IT OUT, AND FOLD IT IN HALF. KEEP IT WITH YOU AT ALL TIMES. IT CAN FIT IN A WALLET, PURSE, OR BACKPACK.

If you are printing this document, select "double-sided" from your printer menu.

INDICATION

SUBLOCADE (buprenorphine extended-release) injection, for subcutaneous use, CIII is a prescription medicine used to treat adults with moderate to severe addiction (dependence) to opioid drugs (prescription or illegal) who have received an oral transmucosal (used under the tongue or inside the cheek) buprenorphine-containing medicine at a dose that controls withdrawal symptoms for at least 7 days. SUBLOCADE is part of a complete treatment plan that should include counseling.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about SUBLOCADE?

Because of the serious risk of potential harm or death from self-injecting SUBLOCADE into a vein (intravenously), it is only available through a restricted program called the SUBLOCADE REMS Program.

- **SUBLOCADE is not available in retail pharmacies.**
- **Your SUBLOCADE injection will only be given to you by a certified healthcare provider.**

SUBLOCADE contains a medicine called buprenorphine. Buprenorphine is an opioid that can cause serious and life-threatening breathing problems, especially if you take or use certain other medicines or drugs.

Talk to your healthcare provider about naloxone. Naloxone is a medicine that is available to patients for the emergency treatment of an opioid overdose. If naloxone is given, you must call 911 or get emergency medical help right away to treat overdose or accidental use of an opioid.

SUBLOCADE may cause serious and life-threatening breathing problems. Get emergency help right away if you:

- feel faint
- feel dizzy
- are confused
- feel sleepy or uncoordinated
- have blurred vision
- have slurred speech
- are breathing slower than normal
- cannot think well or clearly

IMPORTANT SAFETY INFORMATION CONTINUES ON REVERSE PAGE
Please see SUBLOCADE [Full Prescribing Information](#) including **BOXED WARNING and **Medication Guide** at www.sublocade.com.**

P-BAG-US-01501 **EXPIRY AUGUST 2026**

PATIENT ALERT CARD	
In the event that you require medical attention, show this card to a healthcare provider.	
My Name _____	
My Emergency Contact _____	Number _____
My Primary Care Provider _____	Number _____
My Treatment Provider _____	Number _____
SUBLOCADE® (buprenorphine extended-release)	
© 2024 Indivior UK Limited SUBLOCADE® and INDIVIOR® are registered trademarks of Indivior UK Limited All Rights Reserved. P-BAG-US-01501 Intended for US audiences.	
THIS PATIENT IS BEING TREATED WITH SUBLOCADE (buprenorphine extended-release), CIII, A ONCE-MONTHLY SUBCUTANEOUS INJECTION.	
• If a situation arises in which this patient needs acute pain management, treat with a non-opioid analgesic whenever possible.	
• Closely monitor patients who require opioid therapy for analgesia, with particular attention to respiratory function. Higher doses of opioids may be needed for analgesic effect. Therefore, a higher potential for toxicity exists with opioid administration.	
See reverse for BOXED WARNING and see the full Prescribing Information including BOXED WARNING , and Medication Guide at www.sublocade.com .	

ONCE-MONTHLY

Sublocade®
(buprenorphine extended-release)
injection for subcutaneous use
100mg-300mg

IMPORTANT SAFETY INFORMATION (CONT.)

Do not take certain medicines during treatment with SUBLOCADE. Taking other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) while on SUBLOCADE can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.

In an emergency, have family members tell emergency department staff that you are physically dependent on an opioid and are being treated with SUBLOCADE.

You may have detectable levels of SUBLOCADE in your body for a long period after stopping treatment with SUBLOCADE.

Death has been reported in those who are not opioid dependent who received buprenorphine sublingually.

Who should not take SUBLOCADE?

Do not use SUBLOCADE if you are allergic to buprenorphine or any ingredient in the prefilled syringe (Indivior's proprietary buprenorphine gel depot delivery system, a biodegradable 50:50 poly(DL-lactide-co-glycolide) polymer and a biocompatible solvent, N-methyl-2-pyrrolidone (NMP)).

Before starting SUBLOCADE, tell your healthcare provider about all of your medical conditions, including if you have:

- trouble breathing or lung problems
- a curve in your spine that affects your breathing
- Addison's disease
- an enlarged prostate (men)
- problems urinating
- liver, kidney, or gallbladder problems
- alcoholism
- a head injury or brain problem
- mental health problems
- adrenal gland or thyroid gland problems

PATIENT ALERT CARD

Because of the serious risk of potential harm or death from self-injecting SUBLOCADE into a vein (intravenously), it is only available through a restricted program called the SUBLOCADE REMS Program.

- SUBLOCADE is not available in retail pharmacies.
- Your SUBLOCADE injection will only be given to you by a certified healthcare provider.

See reverse and see the full Prescribing Information including **BOXED WARNING**, and Medication Guide at www.sublocade.com.

SUBLOCADE® (buprenorphine extended-release)

This patient is being treated with SUBLOCADE (buprenorphine extended-release) CIII, 100 mg/300 mg, a once-monthly subcutaneous injection.

- If a situation arises in which this patient needs acute pain management, treat with a non-opioid analgesic whenever possible.
- Closely monitor patients who require opioid therapy for analgesia, with particular attention to respiratory function. Higher doses of opioids may be needed for analgesic effect. Therefore, a higher potential for toxicity exists with opioid administration.

PATIENT ALERT CARD

Tell your healthcare provider if you are:

- **pregnant or plan to become pregnant.** If you receive SUBLOCADE while pregnant, your baby may have symptoms of opioid withdrawal at birth that could be life-threatening if not recognized and treated. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- **breastfeeding or plan to breastfeed.** SUBLOCADE can pass into your breast milk and harm your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with SUBLOCADE. Monitor your baby for increased drowsiness and breathing problems if you breastfeed during treatment with SUBLOCADE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

What should I avoid while being treated with SUBLOCADE?

- **Do not drive, operate heavy machinery, or perform any other dangerous activities until you know how SUBLOCADE affects you.** SUBLOCADE can make you sleepy, dizzy, or lightheaded. This may happen more often in the first few days after your injection and when your dose is changed.
- **Do not drink alcohol** or take prescription or over-the-counter medicines that contain alcohol during treatment with SUBLOCADE, because this can lead to loss of consciousness or even death.

IMPORTANT SAFETY INFORMATION CONTINUES ON NEXT PAGE

Please see SUBLOCADE full Prescribing Information including BOXED WARNING and Medication Guide at www.sublocade.com.

P-BAG-US-01501 **EXPIRY AUGUST 2026**

ONCE-MONTHLY

Sublocade®
(buprenorphine extended-release)
injection for subcutaneous use ©
100mg-300mg

IMPORTANT SAFETY INFORMATION (CONT.)

What are the possible side effects of SUBLOCADE?

SUBLOCADE can cause serious side effects, including:

- **Trouble breathing.** Taking other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants during treatment with SUBLOCADE can cause breathing problems that can lead to coma and death.
- **Sleepiness, dizziness, and problems with coordination.**
- **Physical dependence.**
- **Liver problems.** Call your healthcare provider right away if you notice any of these symptoms:
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or “tea-colored” urine
 - light colored stools (bowel movements)
 - loss of appetite
 - pain, aching, or tenderness on the right side of your stomach area
 - nausea
- Your healthcare provider should do blood tests to check your liver before you start and during treatment with SUBLOCADE.
- **Allergic reaction.** You may have rash, hives, itching, swelling of your face, wheezing, low blood pressure, or loss of consciousness. Call your healthcare provider or get emergency help right away.
- **Opioid withdrawal.** Call your healthcare provider right away if you get any of these symptoms:
 - shaking
 - sweating more than normal
 - feeling hot or cold more than normal
 - runny nose
 - watery eyes
 - goose bumps
 - diarrhea
 - vomiting
 - muscle aches
- **Decrease in blood pressure.** You may feel dizzy when you get up from sitting or lying down.
- **The most common side effects of SUBLOCADE include:**
 - constipation
 - headache
 - nausea
 - injection site itching
 - vomiting
 - increase in liver enzymes
 - tiredness
 - injection site pain
- SUBLOCADE may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects. Call your healthcare provider for medical advice about side effects.

To report a pregnancy or side effects associated with taking SUBLOCADE or any safety related information, product complaint, request for medical information, or product query, please contact PatientSafetyNA@indivior.com or 1-877-782-6966. You are encouraged to report negative side effects of drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

See full Prescribing Information, including BOXED WARNING, and Medication Guide, or for more information about SUBLOCADE, visit www.sublocade.com. For REMS information visit www.sublocadeREMS.com. Please see Important Safety Information throughout.

SUBLOCADE is manufactured for Indivior Inc., North Chesterfield VA, 23235.
© 2024 Indivior UK Limited | SUBLOCADE® and INDIVIOR® are registered trademarks of Indivior UK Limited | All Rights Reserved.
P-BAG-US-01501 **EXPIRY AUGUST 2026**
Intended for US audiences.

