

CONTINUING YOUR TREATMENT JOURNEY

SUBLOCADE® (buprenorphine extended-release) injection, for subcutaneous use (CIII) is a prescription medicine used to treat adults with moderate to severe addiction (dependence) to opioid drugs (prescription or illegal) who have received an oral transmucosal (used under the tongue or inside the cheek) buprenorphine-containing medicine at a dose that controls withdrawal symptoms for at least 7 days. SUBLOCADE is part of a complete treatment plan that should include counseling.

Below, You Will Find Information On:

- **The chronic nature of opioid addiction**
- **Tips for maintaining an action plan**
- **Suggestions to help you stay motivated along the way**

A Chronic Condition

Opioid addiction may affect the parts of the brain that control impulses, judgment, and decision-making. For some people with opioid addiction, choosing whether to take opioids can be out of their control.

While opioid addiction is chronic, it can be treatable. It's important to follow your complete treatment plan.

Ask your healthcare provider about the length of your treatment with SUBLOCADE.

Continuing the Conversation With a Healthcare Provider

You can work towards your goals by talking with your healthcare provider team to create a plan of action, which should include counseling. Your treatment team can help with questions or concerns you may have along the way.

Here are some questions you may want to discuss with your treatment team:

1. What's a strategy for staying on track with my treatment plan?

2. How do I start the conversation with my loved ones about my condition?

IMPORTANT SAFETY INFORMATION

Because of the serious risk of potential harm or death from self-injecting SUBLOCADE into a vein (intravenously), it is only available through a restricted program called the SUBLOCADE REMS Program.

- SUBLOCADE is not available in retail pharmacies.
- Your SUBLOCADE injection will only be given to you by a certified healthcare provider.

Please see SUBLOCADE Prescribing Information including **BOXED WARNING** and **Medication Guide** at sublocade.com and full Important Safety Information on pages 7–8.

3. What are some things I can do to structure my day?

Do not stop treatment abruptly. Talk to your healthcare provider first.

Jot It Down

Write down anything you want to share with your healthcare provider or counselor. Examples can include your goals and things that cause stress.

People You Can Count On

Do you have people you can speak to when you need help? You can list people who may help you throughout your journey.

IMPORTANT SAFETY INFORMATION (CONT'D)

In an emergency, you or your family should tell the emergency medical staff that you are physically dependent on an opioid and are being treated with SUBLOCADE.

Buprenorphine, the medicine in SUBLOCADE, can cause serious and life-threatening problems, especially if you take or use certain other medicines or drugs. Call your healthcare provider right away or get emergency help if you: feel faint or dizzy, have mental changes such as confusion, have slower breathing than you normally have, have severe sleepiness, have blurred vision, have problems with coordination, have slurred speech, cannot think well or clearly, have a high body temperature, have slowed reflexes, feel agitated, have stiff muscles, or have trouble walking. These can be signs of an overdose or other serious problems.


Death or serious harm, including life-threatening breathing problems, can happen if you take anxiety medicines or benzodiazepines, sleeping pills, tranquilizers, muscle relaxants, or sedatives, antidepressants, or antihistamines, or drink alcohol during treatment with SUBLOCADE. Tell your healthcare provider if you are taking any of these medicines and if you drink alcohol.

SUBLOCADE is a controlled substance (CIII) because it contains buprenorphine that can be a target for people who abuse prescription medicines or street drugs.

Death has been reported in those who are not opioid dependent who received buprenorphine sublingually.

Please see SUBLOCADE Prescribing Information including BOXED WARNING and Medication Guide at sublocade.com and full Important Safety Information on pages 7–8.

P-BAG-US-00735



Keeping Track of Your Appointments

To stay on track with your treatment plan, it's important to schedule and keep up with your appointments. Schedule them in advance so that your SUBLOCADE prescription is ready at your visit. You can use the space provided to help remember appointments and things that may be discussed during your visits.

Write Down the Dates

Initial appointment

Follow-up appointments

Counseling appointments

IMPORTANT SAFETY INFORMATION (CONT'D)

Do not use SUBLOCADE if you are allergic to buprenorphine or any ingredient in the prefilled syringe (ATRIGEL® Delivery System, a biodegradable 50:50 poly(DL-lactide-co-glycolide) polymer and a biocompatible solvent, *N*-methyl-2-pyrrolidone (NMP)).

SUBLOCADE may not be right for you. Before starting SUBLOCADE, tell your healthcare provider about all of your medical conditions, including: trouble breathing or lung problems, an enlarged prostate gland (men), a head injury or brain problem, problems urinating, a curve in your spine that affects your breathing (scoliosis), liver problems, gallbladder problems, adrenal gland problems, Addison's disease, low thyroid hormone levels (hypothyroidism), a history of alcoholism, mental problems such as hallucinations (seeing or hearing things that are not there), or if you are pregnant or plan to become pregnant. Opioid-dependent women on buprenorphine maintenance therapy may require additional analgesia during labor. If you receive SUBLOCADE while pregnant, your baby may have symptoms of opioid withdrawal at birth. If you are breastfeeding or plan to breastfeed, SUBLOCADE can pass into your breast milk and may harm your baby. Talk with your healthcare provider about the best way to feed your baby during treatment with SUBLOCADE. Watch your baby for increased drowsiness and breathing problems.

Please see SUBLOCADE Prescribing Information including **BOXED WARNING** and **Medication Guide** at sublocade.com and full Important Safety Information on pages 7–8.

P-BAG-US-00735

ONCE-MONTHLY

Sublocade[®]
(buprenorphine extended-release)
injection for subcutaneous use [®]
100mg-300mg

Questions you may want to ask your healthcare provider or counselor

Additional information you may want to remember from your appointments or counseling sessions

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. SUBLOCADE may affect the way other medicines work and other medicines may affect how SUBLOCADE works. Some medicines may cause serious or life-threatening medical problems when taken with SUBLOCADE. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.


The doses of certain medicines may need to be changed if used during treatment with SUBLOCADE. Do not take any medicine during treatment with SUBLOCADE until you have talked with your healthcare provider. Your healthcare provider will tell you if it is safe to take other medicines during treatment with SUBLOCADE.

You should not take anxiety medicines or benzodiazepines (such as Valium® or Xanax®), sleeping pills, tranquilizers, muscle relaxants, or sedatives (such as Ambien®), antidepressants, or antihistamines that are not prescribed to you during treatment with SUBLOCADE, as this can lead to slowed breathing, drowsiness, delayed reaction time, loss of consciousness or even death. If a healthcare provider is considering prescribing such a medicine for you, remind the healthcare provider that you are being treated with SUBLOCADE.

You may have detectable levels of SUBLOCADE in your body for a long period after stopping treatment with SUBLOCADE.

Please see SUBLOCADE Prescribing Information including BOXED WARNING and Medication Guide at sublocade.com and full Important Safety Information on pages 7–8.

P-BAG-US-00735



Staying Motivated and Remembering Why You Started Treatment

As you progress, you may also experience setbacks. Remember why you are seeking treatment, and talk to your healthcare provider along the way.

You can reach out to your treatment team or support group to help keep you motivated throughout your journey.

The provided space can also be used to create a checklist as a reminder of why you started treatment, when you need additional motivation to stay on course. Once you've completed this list, you can refer back to it to see the goals you have set along the way.

Write the Reasons You Chose to Seek Treatment

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I avoid while being treated with SUBLOCADE?

- **Do not drive, operate heavy machinery, or perform any other dangerous activities until you know how this medicine affects you.** Buprenorphine can cause drowsiness and slow reaction times. This may happen more often in the first few days after your injection and when your dose is changed.
- **Do not drink alcohol** during treatment with SUBLOCADE, as this can lead to slowed breathing, drowsiness, slow reaction time, loss of consciousness or even death.


What are the possible side effects of SUBLOCADE?

SUBLOCADE can cause serious side effects, including:

- **Physical dependence and withdrawal.** Your body can develop a physical need for SUBLOCADE (dependence). If you stop receiving SUBLOCADE, you could have opioid withdrawal symptoms such as: shaking, goose bumps, muscle aches, sweating more than normal, feeling hot or cold more than normal, runny nose and watery eyes, or diarrhea or vomiting. These symptoms may start weeks to months after your last dose of SUBLOCADE.

Please see SUBLOCADE Prescribing Information including **BOXED WARNING** and **Medication Guide** at sublocade.com and full Important Safety Information on pages 7–8.

P-BAG-US-00735



Remember Your Milestones

IMPORTANT SAFETY INFORMATION (CONT'D)

- **Liver problems.** Call your healthcare provider right away if you notice any of these signs of liver problems: your skin or the white part of your eyes turns yellow (jaundice), urine turns dark, bowel movements (stools) turn light in color, decreased appetite, or stomach (abdomen) pain or nausea. Your healthcare provider may do tests before and during treatment with SUBLOCADE to check your liver.
- **Allergic reaction.** Call your healthcare provider or get emergency help right away if you get: rash, hives, itching, swelling of your face, wheezing or dizziness, or a decrease in consciousness.
- **Decrease in blood pressure.** You may feel dizzy when you get up from sitting or lying down.
- **The most common side effects of SUBLOCADE include:** constipation, headache, nausea, injection site itching, vomiting, increase in liver enzymes, tiredness, or injection site pain.
- Long-term (chronic) use of opioids, including SUBLOCADE, may cause fertility problems in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects. Call your healthcare provider for medical advice about side effects.

To report pregnancy or side effects associated with taking SUBLOCADE, please call 1-877-782-6966. You are encouraged to report negative side effects of drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For more information about SUBLOCADE, the full Prescribing Information including BOXED WARNING, and Medication Guide visit www.sublocade.com or talk to your healthcare provider. For REMS information visit www.sublocadeREMS.com.

Please see SUBLOCADE Prescribing Information including **BOXED WARNING** and **Medication Guide** at sublocade.com and full Important Safety Information on pages 7–8.

P-BAG-US-00735

ONCE-MONTHLY

Sublocade[®]
(buprenorphine extended-release)
injection for subcutaneous use [®]
100mg•300mg

INDICATION & IMPORTANT SAFETY INFORMATION

INDICATION

SUBLOCADE® (buprenorphine extended-release) injection, for subcutaneous use (CIII) is a prescription medicine used to treat adults with moderate to severe addiction (dependence) to opioid drugs (prescription or illegal) who have received an oral transmucosal (used under the tongue or inside the cheek) buprenorphine-containing medicine at a dose that controls withdrawal symptoms for at least 7 days. SUBLOCADE is part of a complete treatment plan that should include counseling.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about SUBLOCADE?

Because of the serious risk of potential harm or death from self-injecting SUBLOCADE into a vein (intravenously), it is only available through a restricted program called the SUBLOCADE REMS Program.

- SUBLOCADE is not available in retail pharmacies.
- Your SUBLOCADE injection will only be given to you by a certified healthcare provider.

In an emergency, you or your family should tell the emergency medical staff that you are physically dependent on an opioid and are being treated with SUBLOCADE.

Buprenorphine, the medicine in SUBLOCADE, can cause serious and life-threatening problems, especially if you take or use certain other medicines or drugs. Call your healthcare provider right away or get emergency help if you: feel faint or dizzy, have mental changes such as confusion, have slower breathing than you normally have, have severe sleepiness, have blurred vision, have problems with coordination, have slurred speech, cannot think well or clearly, have a high body temperature, have slowed reflexes, feel agitated, have stiff muscles, or have trouble walking. These can be signs of an overdose or other serious problems.

Death or serious harm, including life-threatening breathing problems, can happen if you take anxiety medicines or benzodiazepines, sleeping pills, tranquilizers, muscle relaxants, or sedatives, antidepressants, or antihistamines, or drink alcohol during treatment with SUBLOCADE. Tell your healthcare provider if you are taking any of these medicines and if you drink alcohol.

SUBLOCADE is a controlled substance (CIII) because it contains buprenorphine that can be a target for people who abuse prescription medicines or street drugs.

Death has been reported in those who are not opioid dependent who received buprenorphine sublingually.

Do not use SUBLOCADE if you are allergic to buprenorphine or any ingredient in the prefilled syringe (ATRIGEL® Delivery System, a biodegradable 50:50 poly(DL-lactide-co-glycolide) polymer and a biocompatible solvent, *N*-methyl-2-pyrrolidone (NMP)).

SUBLOCADE may not be right for you. Before starting SUBLOCADE, tell your healthcare provider about all of your medical conditions, including: trouble breathing or lung problems, an enlarged prostate gland (men), a head injury or brain problem, problems urinating, a curve in your spine that affects your breathing (scoliosis), liver problems, gallbladder problems, adrenal gland problems, Addison's disease, low thyroid hormone levels (hypothyroidism), a history of alcoholism, mental problems such as hallucinations (seeing or hearing things that are not there), or if you are pregnant or plan to become pregnant. Opioid-dependent women on buprenorphine maintenance therapy may require additional analgesia during labor. If you receive SUBLOCADE while pregnant, your baby may have symptoms of opioid withdrawal at birth. If you are breastfeeding or plan to breastfeed, SUBLOCADE can pass into your breast milk and may harm your baby. Talk with your healthcare provider about the best way to feed your baby during treatment with SUBLOCADE. Watch your baby for increased drowsiness and breathing problems.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. SUBLOCADE may affect the way other medicines work and other medicines may affect how SUBLOCADE works. Some medicines may cause serious or life-threatening medical problems when taken with SUBLOCADE. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

The doses of certain medicines may need to be changed if used during treatment with SUBLOCADE. Do not take any medicine during treatment with SUBLOCADE until you have talked with your healthcare provider. Your healthcare provider will tell you if it is safe to take other medicines during treatment with SUBLOCADE.

You should not take anxiety medicines or benzodiazepines (such as Valium® or Xanax®), sleeping pills, tranquilizers, muscle relaxants, or sedatives (such as Ambien®), antidepressants, or antihistamines that are not prescribed to you during treatment with SUBLOCADE, as this can lead to slowed breathing, drowsiness, delayed reaction time, loss of consciousness or even death. If a healthcare provider is considering prescribing such a medicine for you, remind the healthcare provider that you are being treated with SUBLOCADE.

You may have detectable levels of SUBLOCADE in your body for a long period after stopping treatment with SUBLOCADE.

Please see SUBLOCADE Prescribing Information including **BOXED WARNING** and Medication Guide at sublocade.com and full Important Safety Information continued on page 8.

P-BAG-US-00735

ONCE-MONTHLY

Sublocade[®]
(buprenorphine extended-release)
injection for subcutaneous use 
100mg-300mg

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I avoid while being treated with SUBLOCADE?

- **Do not drive, operate heavy machinery, or perform any other dangerous activities until you know how this medicine affects you.** Buprenorphine can cause drowsiness and slow reaction times. This may happen more often in the first few days after your injection and when your dose is changed.
- **Do not drink alcohol** during treatment with SUBLOCADE, as this can lead to slowed breathing, drowsiness, slow reaction time, loss of consciousness or even death.

What are the possible side effects of SUBLOCADE?

SUBLOCADE can cause serious side effects, including:

- **Physical dependence and withdrawal.** Your body can develop a physical need for SUBLOCADE (dependence). If you stop receiving SUBLOCADE, you could have opioid withdrawal symptoms such as: shaking, goose bumps, muscle aches, sweating more than normal, feeling hot or cold more than normal, runny nose and watery eyes, or diarrhea or vomiting. These symptoms may start weeks to months after your last dose of SUBLOCADE.
- **Liver problems.** Call your healthcare provider right away if you notice any of these signs of liver problems: your skin or the white part of your eyes turns yellow (jaundice), urine turns dark, bowel movements (stools) turn light in color, decreased appetite, or stomach (abdomen) pain or nausea. Your healthcare provider may do tests before and during treatment with SUBLOCADE to check your liver.
- **Allergic reaction.** Call your healthcare provider or get emergency help right away if you get: rash, hives, itching, swelling of your face, wheezing or dizziness, or a decrease in consciousness.
- **Decrease in blood pressure.** You may feel dizzy when you get up from sitting or lying down.
- **The most common side effects of SUBLOCADE include:** constipation, headache, nausea, injection site itching, vomiting, increase in liver enzymes, tiredness, or injection site pain.
- Long-term (chronic) use of opioids, including SUBLOCADE, may cause fertility problems in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects. Call your healthcare provider for medical advice about side effects.

To report pregnancy or side effects associated with taking SUBLOCADE, please call 1-877-782-6966. You are encouraged to report negative side effects of drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For more information about SUBLOCADE, the full Prescribing Information including BOXED WARNING, and Medication Guide visit www.sublocade.com or talk to your healthcare provider. For REMS information visit www.sublocadeREMS.com.

SUBLOCADE is a registered trademark of Indivior UK Limited.
© Indivior PLC | INDIVIOR is a registered trademark of Indivior UK Limited.
All rights reserved. SUBLOCADE is manufactured for Indivior Inc.,
North Chesterfield, VA 23235 by AMRI, Burlington, MA 01803.
P-BAG-US-00735 EXPIRY AUGUST 2022

This material is intended solely for residents of the United States.



INDIVIOR™

ONCE-MONTHLY

Sublocade®

(buprenorphine extended-release)
injection for subcutaneous use Ⓢ
100mg•300mg